

ENDOMORPHISMS OF THE PLANE PRESERVING A PENCIL OF CURVES

MARIUS DABIJA AND MATTIAS JONSSON

ABSTRACT. We classify endomorphisms of the plane that preserve a pencil of curves.

INTRODUCTION

In dynamical systems, an interesting role is played by systems that are *integrable* in the sense that they preserve some additional structure. In this note we are concerned with endomorphisms of the complex projective plane $\mathbf{P}^2$ that preserve a pencil of curves.¹

More precisely, let $f : \mathbf{P}^2 \rightarrow \mathbf{P}^2$ be an endomorphism of algebraic degree $d > 1$ and let $\mathcal{P}$ be a pencil of curves of degree $k \geq 1$ on $\mathbf{P}^2$, defined by $\pi : \mathbf{P}^2 \dashrightarrow \mathbf{P}^1$. For simplicity we shall assume that $\mathcal{P}$ is *irreducible*, that is, the generic curve in $\mathcal{P}$ is irreducible. (The reducible case is discussed in Section 3.) We say that f leaves $\mathcal{P}$ *invariant* if f maps any curve in $\mathcal{P}$ (on)to another curve in $\mathcal{P}$. This implies that there exists an endomorphism $g : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ of degree d such that $\pi \circ f = g \circ \pi$.

Main Theorem. *There are only two types of invariant irreducible pencils $\mathcal{P}$.*

- (i) $\mathcal{P}$ is an elementary pencil, that is, the pencil of lines through a point. Pick homogeneous coordinates on $\mathbf{P}^2$ and $\mathbf{P}^1$ such that $\pi[x : y : z] = [x : y]$. Then f takes the form $f[x : y : z] = [P(x, y) : Q(x, y) : R(x, y, z)]$.
- (ii) $\mathcal{P}$ is a binomial pencil. This means that we may pick homogeneous coordinates on $\mathbf{P}^2$ and $\mathbf{P}^1$ such that $\pi[x : y : z] = [x^h y^{k-h} : z^k]$ where $0 < h < k$ and $\gcd(h, k) = 1$. In this case, after conjugation with an element of $\mathrm{PGL}(3, \mathbf{C})$ preserving $\mathcal{P}$, f takes the form $f[x : y : z] = [x^d : y^d : R]$ or $f[x : y : z] = [y^d : x^d : R]$ for $R(x, y, z) = z^{d-kl} \prod_{i=1}^l (z^k + c_i x^h y^{k-h})$, where $0 \leq l \leq d/k$ and $c_i \in \mathbf{C}^*$.

We shall only prove that an invariant (irreducible) pencil $\mathcal{P}$ must be elementary or binomial. The verification that f is of the form stated in the theorem is straightforward and left to the reader.

Date: February 2, 2008.

Marius Dabija passed away on June 22, 2003.

¹See [Si] for a survey of dynamics on projective spaces and [J1, J2] for dynamics of endomorphisms appearing in the Main Theorem.

Our main theorem should be compared to the classification of curves in $\mathbf{P}^2$ that are totally invariant by an endomorphism. Such a curve must be a union of at most three lines [FS, D, CL, BCS], a result that we exploit in the proof.

The strategy is to study the interplay between the ramification loci of π , f and g . In doing so, the reducible and multiple curves in $\mathcal{P}$ play an important role. We also use the minimal desingularization of $\mathcal{P}$ and the Riemann-Hurwitz formula to analyze the induced map between fibers.

1. COUNTING RAMIFICATION DEGREES

Let $\mathcal{P}$ be an irreducible pencil of plane curves of degree k with associated projection $\pi : \mathbf{P}^2 \dashrightarrow \mathbf{P}^1$, say $\pi[x : y : z] = [A(x, y, z) : B(x, y, z)]$ in homogeneous coordinates. By an *element* of $\mathcal{P}$ we shall mean the divisor of $sA + tB$ for some $[s : t] \in \mathbf{P}^1$. A *curve* in $\mathcal{P}$ will mean the (reduced) support of an element of $\mathcal{P}$.

The *base locus* $\mathcal{B}(\mathcal{P})$ is the (nonempty, finite) indeterminacy set of π . The *ramification locus* R_π is defined as follows. For an effective divisor $D = \sum_j m_j D_j$ on $\mathbf{P}^2$, set $R_D = \sum_j (m_j - 1) D_j$. Then $R_\pi = \sum R_D$, the sum being over the elements of $\mathcal{P}$. Set $e(\mathcal{P}) := 2k - \deg R_\pi$.

Lemma 1.1. *We have $e(\mathcal{P}) \geq 2$ with equality iff $\mathcal{P}$ is an elementary pencil.*

Proof. Consider a generic line $L \subset \mathbf{P}^2$. Then $L \cap \mathcal{B}(\mathcal{P}) = \emptyset$, L is transversal to R_π and all tangencies (if any) between L and the curves in $\mathcal{P}$ are quadratic. Applying the Riemann-Hurwitz formula to $\pi|_L : L \rightarrow \mathbf{P}^1$, which has topological degree k , we obtain $2 = 2k - \deg R_\pi - t$, where $t \geq 0$ is the number of tangencies as above. In particular, $t = 0$ iff there are no tangencies and this is easily seen to happen iff $\mathcal{P}$ is elementary. Since $e(\mathcal{P}) = 2 + t$, the proof is complete. $\square$

Now assume that $f : \mathbf{P}^2 \rightarrow \mathbf{P}^2$ is an endomorphism preserving $\mathcal{P}$. Let $g : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ be the induced endomorphism, that is, $\pi \circ f = g \circ \pi$. Denote the ramification loci of f and g by $R_f \subset \mathbf{P}^2$ and $R_g \subset \mathbf{P}^1$, respectively. Write $R_f^\mathcal{P}$ for the part of R_f supported on curves in $\mathcal{P}$. A direct calculation based on $\pi \circ f = g \circ \pi$ yields

Lemma 1.2. *We have $R_f^\mathcal{P} + f^* R_\pi = R_\pi + \pi^* R_g$. In particular, $\deg R_f^\mathcal{P} = e(d - 1)$, with $e = e(\mathcal{P})$ as above.*

Since $\deg R_f = 3(d - 1)$ and $e \geq 2$ we see that $e = 2$ or $e = 3$. If $e = 2$, then $\mathcal{P}$ is an elementary pencil by Lemma 1.1, so let us assume from now on that $e = 3$. In particular, R_f is supported on curves in $\mathcal{P}$.

Lemma 1.3. *Assume $\mathcal{P}$ has a reducible curve. Then $\mathcal{P}$ is a binomial pencil.*

Proof. As f maps curves in $\mathcal{P}$ onto curves in $\mathcal{P}$, the reducible curves in $\mathcal{P}$ must be totally invariant for f . Since any totally invariant curve for f is a union of at most three lines [FS, D, CL, BCS], there is exactly one reducible curve, the union of two or three lines.

In the case of two lines, we assume $A(x, y, z) = x^h y^{k-h}$ with $0 < h < k$. Let $m_j C_j$ be the multiple curves of $\mathcal{P}$, i.e. $m_j \geq 2$ and C_j is reduced and irreducible. Then $R_\pi = \sum_{j=1}^J (m_j - 1) C_j + (h - 1)\{x = 0\} + (k - h - 1)\{y = 0\}$. Taking degrees and using $\deg R_\pi = 2k - 3$ we get $\sum_{j=1}^J (1 - \frac{1}{m_j}) = 1 - \frac{1}{k}$. Since $k > 1$, the unique solution is given by $J = 1$, $m_1 = k$. Thus $\mathcal{P}$ admits a multiple line. Since $\mathcal{P}$ is irreducible, this line cannot pass through $[0 : 0 : 1]$, so we may assume $B(x, y, z) = z^k$. Then $\mathcal{P}$ is a binomial pencil.

When the reducible curve is a union of three lines we may assume $A(x, y, z) = x^h y^l z^{k-h-l}$ with $h, l > 0$, $h + l < k$. Reasoning as above we get $\sum_{j=1}^J (1 - \frac{1}{m_j}) = 1$, implying $J = 2$ and $m_1 = m_2 = 2$. But then $\mathcal{P}$ is reducible, a contradiction. $\square$

2. RULING OUT THE REMAINING CASES

From now on we assume $e = e(\mathcal{P}) = 3$, that $\mathcal{P}$ has no reducible curve, and that f preserves $\mathcal{P}$. In the end, we shall see that this is impossible.

Let $p : X \rightarrow \mathbf{P}^2$ be a desingularization of the pencil $\mathcal{P}$, i.e. p is a composition of point blowups and $\pi : \mathbf{P}^2 \dashrightarrow \mathbf{P}^1$ lifts to a regular map $\pi_X : X \rightarrow \mathbf{P}^1$ defining a ruling on X . Let $\text{Exc}(p) \subset X$ be the exceptional locus of p ; we call its irreducible components *exceptional primes*. There are finitely many exceptional primes E_i , $i = 1, \dots, I$ of $\text{Exc}(p)$ that are *horizontal*, i.e. π_X maps E_i surjectively onto $\mathbf{P}^1$, say with degree n_i . Write E' for the union of the other (vertical) exceptional primes. Let $F_t = \pi_X^{-1}(t)$, $t \in \mathbf{P}^1$, be the fibers of π_X . For generic t , F_t is smooth, intersects E_i transversely at $n_i \geq 1$ distinct generic points, and is disjoint from E' .

Recall that R_f is supported on curves in $\mathcal{P}$. Moreover, since f is finite, the base locus $\mathcal{B}(\mathcal{P}) \subset \mathbf{P}^2$ is totally invariant. Thus, for generic t , f induces a regular map of F_t onto $F_{g(t)}$ and restricts to a covering map of $F_t \setminus \text{Exc}(p)$ onto $F_{g(t)} \setminus \text{Exc}(p)$. By Riemann-Hurwitz this implies that F_t is rational and that $\sum_{i=1}^I n_i = 2$. In particular $\mathcal{P}$ has one or two base points in $\mathbf{P}^2$.

As before, let $m_j C_j$ be the multiple curves of $\mathcal{P}$. Since $\mathcal{P}$ has no reducible curves, the ramification locus R_π is supported on the C_j 's. Counting degrees as above, we get $2 - \frac{3}{k} = \sum_{j=1}^J (1 - \frac{1}{m_j})$. As $\mathcal{P}$ is irreducible, the multiplicities m_j are pairwise relatively prime. One easily checks that the only two possibilities are $k = 2, J = 1, m_1 = 2$ and $k = 90, J = 3, m_1 = 2, m_2 = 3, m_3 = 5$, respectively.

In the first case we have a pencil of conics with a single double line. When there are two base points in $\mathbf{P}^2$, one checks that $\mathcal{P}$ has a reducible curve (and is binomial), a contradiction. If there is a single base point in $\mathbf{P}^2$, we may assume $A = y + x^2$, $B = y^2$. Then $I = 1$, $n_1 = 1$ in the notation above, contradicting $\sum n_i = 2$.

In the second case, we have three multiple curves, of multiplicities 2, 3 and 5, respectively. We may and will assume that the resolution $p : X \rightarrow \mathbf{P}^2$ of $\mathcal{P}$ is *minimal*, a condition that may be paraphrased as follows. Let ν_i be the divisorial valuation on the function field of $\mathbf{P}^2$ associated to the horizontal exceptional prime

E_i , $1 \leq i \leq I$. Then $p : X \rightarrow \mathbf{P}^2$ is minimal with the property that the centers of ν_i on X are all one-dimensional. By considering the dual graph of p from this point of view (see e.g. [Sp, FJ]) one finds that at least one of the horizontal exceptional primes, say E_1 , intersects the other exceptional primes in a finite set Y consisting of at most *two* points. Thus at least one of the multiple fibers $F_t = \pi_X^{-1}(t)$, say F_0 , must intersect E_1 but not Y . As noted above, a generic fiber F_t intersects E_1 transversely at n_1 distinct points. Hence $(F_t \cdot E_1) = n_1$ and this equation must then hold for *all* t . In particular it holds for the multiple fiber F_0 . As $\sum n_i = 2$, this is only possible if $I = 1$, $n_1 = 2$ and $F_0 = 2G_0$ is a double curve. Here G_0 is a priori reducible, but intersects E_1 transversely at a single point, not in Y .

Now pick a generic fiber F_t and suppose that $g(t) = 0$, that is, the lift of f to X maps F_t into G_0 . Then the two intersection points of F_t and E_1 must map to the unique intersection point of G_0 and E_1 . Thus G_0 must be irreducible and we get a covering map of degree $d > 1$ from $F_t \setminus E_1$ onto $G_0 \setminus E_1$. By Riemann-Hurwitz, this is impossible.

Hence, for t generic we cannot have $g(t) = 0$ and in fact not $g^n(t) = 0$ either for any $n \geq 1$. This implies that $t = 0$ is totally invariant for g^2 , so the corresponding curve C_0 in $\mathbf{P}^2$ is totally invariant for f^2 , a contradiction since $\deg C_0 = 45 > 3$.

This concludes the proof of the Main Theorem.

3. REDUCIBLE PENCILS AND FURTHER REMARKS

Finally we discuss the case when $\mathcal{P}$ is reducible (but without base divisor). One must then be a little careful with the notion of invariance. For example, let $\mathcal{P}$ be defined by $\pi[x : y : z] = [x^2 : y^2]$ and set $f[x : y : z] = [x^2 : y^2 : z^2]$. Then f maps curves in $\mathcal{P}$ into, but not onto, curves in $\mathcal{P}$. On the other hand, the preimage of a fiber is a union of fibers and we shall use this as the definition of invariance.

It is a result of Bertini that $\mathcal{P}$ factors through an irreducible pencil $\mathcal{P}'$, say defined by $\pi' : \mathbf{P}^2 \dashrightarrow \mathbf{P}^1$. This means that $\pi = \phi \circ \pi'$, where $\phi : \mathbf{P}^1 \rightarrow \mathbf{P}^1$ is an endomorphism of degree > 1 . Clearly, if $\mathcal{P}$ is invariant by f , then so is $\mathcal{P}'$, which implies that $\mathcal{P}'$ is elementary or binomial and that f is as described in the Main Theorem.

To completely solve the classification problem in the reducible case would therefore amount to the following problem. Given an endomorphism g' of $\mathbf{P}^1$ associated to one of the situations in the Main Theorem, find all endomorphisms ϕ of $\mathbf{P}^1$ of degree > 1 such that $\phi \circ g' = g \circ \phi$ for another endomorphism g . This appears to be an interesting and nontrivial problem, but one of a different flavor than the present note, and we shall leave it for now.

Beyond the setup used here, there are numerous possible notions of integrable holomorphic dynamics. One could look at both more general (rational) mappings, more general structures (linear systems, webs, foliations, ...) and of course higher dimensions. In [DJ] we study algebraic webs on $\mathbf{P}^2$ invariant under endomorphisms. Cantat and Favre [CF] have classified infinite groups of birational surface maps

preserving a holomorphic foliation. Pereira [P] has outlined a generalization of the results in this paper from pencils to foliations. However, many interesting problems remain open.

Acknowledgment. The second author was supported by the NSF, the Swedish Research Council and the Gustafsson Foundation. He is grateful to Igor Dolgachev, Charles Favre and Jorge Pereira for valuable discussions. He also thanks the referee for suggesting substantial simplifications to several arguments in a previous version of the paper.

REFERENCES

- [BCS] J.-Y. Briend, S. Cantat and M. Shishikura. *Linearity of the exceptional set for maps of $\mathbf{P}^k(\mathbb{C})$* . Math. Ann., **330** (2004), 39–43.
- [CF] S. Cantat and C. Favre. *Symétries birationnelles des surfaces feuilletées*. J. Reine Angew. Math., **561** (2003), 199–235.
- [CL] D. Cerveau and A. Lins Neto. *Hypersurfaces exceptionnelles des endomorphismes de $\mathbf{C}^n$* . Bol. Soc. Brasil. Mat. (N.S.), **31** (2000), 155–161.
- [D] M. Dabija. *Algebraic and geometric dynamics in several complex variables*. Thesis. Univ. of Michigan, 2000.
- [DJ] M. Dabija and M. Jonsson. *Algebraic webs invariant under endomorphisms*. In preparation.
- [FJ] C. Favre and M. Jonsson. *The valuative tree*. Lecture Notes in Mathematics, 1853, Springer-Verlag, Berlin, 2004.
- [FS] J.-E. Fornæss and N. Sibony. *Complex dynamics in higher dimension, I*. In *Complex analytic methods in dynamical systems (Rio de Janeiro, 1992)*, Astérisque, **222** (1994), 201–231.
- [J1] M. Jonsson. *Dynamics of polynomial skew products on $\mathbf{C}^2$* . Math. Ann. **314** (1999), 403–447.
- [J2] M. Jonsson. *Ergodic theory of fibered rational maps*. Ark. Math. **38** (2000), 281–317.
- [P] J.V. Pereira. Personal communication.
- [Si] N. Sibony. *Dynamique des applications rationnelles de $\mathbf{P}^k$* . In *Dynamique et géométrie complexes (Lyon, 1997)*. Panor. Synthèses, 8 (1999), 97–185.
- [Sp] M. Spivakovsky. *Valuations in function fields of surfaces*. Amer. J. Math. **112** (1990), 107–156.

DEPT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, ANN ARBOR, MI 48109-1109, USA, AND,
 DEPT OF MATHEMATICS, KTH, SE-100 44 STOCKHOLM, SWEDEN
E-mail address: mattiasj@umich.edu, mattiasj@kth.se